

BENEŠ CONDITION FOR DISCONTINUOUS EXPONENTIAL MARTINGALE

R. LIPTSER

ABSTRACT. It is known the Girsanov exponent $\mathfrak{z}_t$, being solution of Doleans-Dade equation $\mathfrak{z}_t = 1 + \int_0^t \alpha(\omega, s) dB_s$ generated by Brownian motion B_t and a random process $\alpha(\omega, t)$ with $\int_0^t \alpha^2(\omega, s) ds < \infty$ a.s., is the martingale provided that the Beneš condition

$$|\alpha(\omega, t)|^2 \leq \text{const.} \left[1 + \sup_{s \in [0, t]} B_s^2 \right], \quad \forall t > 0,$$

holds true. In this paper, we show B_t can be replaced by a homogeneous purely discontinuous square integrable martingale M_t with independent increments and paths from the Skorokhod space $\mathbb{D}_{[0, \infty)}$ having positive jumps ΔM_t with $\mathbb{E} \sum_{s \in [0, t]} (\Delta M_s)^3 < \infty$. A function $\alpha(\omega, t)$ is assumed to be nonnegative and predictable. Under this setting $\mathfrak{z}_t$ is the martingale provided that

$$\alpha^2(\omega, t) \leq \text{const.} \left[1 + \sup_{s \in [0, t]} M_{s-}^2 \right], \quad \forall t > 0.$$

The method of proof differs from the original Beneš one and is compatible for both setting with B_t and M_t .

1. Introduction and main result

1.1. Setting of problem. A classical Girsanov's exponent

$$\mathfrak{z}_t = \exp \left(\int_0^t \alpha(\omega, s) dB_s - \frac{1}{2} \int_0^t \alpha^2(\omega, s) ds \right),$$

with Brownian motion B_t and adapted random process $\alpha(\omega, t)$ having $\int_0^t \alpha(\omega, s) ds < \infty$, forms a positive local martingale (and supermartingale too) with $\mathbb{E} \mathfrak{z}_t \leq 1$. If

$$\mathbb{E} \mathfrak{z}_t \equiv 1, \tag{1.1}$$

the random process $\mathfrak{z}_t$ is a martingale. In order (1.1) to have, Girsanov in [2] used bounded function $\alpha(\omega, t)$ and suggested a conjecture that (1.1) will be valid if $\alpha^2(\omega, t) \approx B_t^2$. Among conditions guaranteing (1.1) (see, e.g. Novikov [10], Kazamaki [6], the latest Krylov [7], etc), we distinguish the Beneš statement: for any $T > 0$,

$$“|\alpha(\omega, t)|^2 \leq \text{const.} \left[1 + \sup_{s \in [0, t]} B_s^2 \right]_{t \in [0, T]}” \Rightarrow “\mathbb{E} \mathfrak{z}_T = 1”, \tag{1.2}$$

which is derived in [1] with the help of Kazamaki [6] (see also Karatzas and Shreve [5], “Ustünel and Zakai [11]).

1991 *Mathematics Subject Classification.* 60G45, 60G46 .

Key words and phrases. Girsanov, exponential martingale, uniform integrability.

The aim of this paper is to “replace” the Brownian motion B_t by a homogeneous purely discontinuous square integrable martingale M_t with independent increments and obtain an implication similar to (1.2). Unfortunately, a method of proof of (1.2) with M_t instead of B_t is not applicable since, for example, results of Kazamaki, Novikov, and Krylov related to martingale M_t do not exist.

1.2. New approach to Beneš result. We propose a new approach for the proof of (1.2) which is compatible with B_t and M_t . Its application to a discontinuous martingale is more involved than to classical case with Brownian motion. So, to make our approach at most transparent we give a sketch of the proof of (1.2) step by step.

(1) It is well known $\mathfrak{z}_t$ is the unique solution of Doléans-Dade equation

$$\mathfrak{z}_t = 1 + \int_0^t \mathfrak{z}_s \alpha(\omega, s) dB_s. \quad (1.3)$$

Set $\sigma_n = \inf \left\{ t : \left[1 + \sup_{s \in [0, t]} B_s^2 \right] \geq n \right\}$, $B_t^n = B_{t \wedge \sigma_n}$ and $\mathfrak{z}_t^n = \mathfrak{z}_{t \wedge \sigma_n}$. Then

$$\mathfrak{z}_t^n = 1 + \int_0^t \mathfrak{z}_s^n I_{\{\sigma_n \geq s\}} \alpha(\omega, s) dB_s.$$

Evidently, $I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) \leq \text{const.} n$. Consequently, $\mathbb{E} \mathfrak{z}_t^n \equiv 1$.

(2) In order to prove $\mathbb{E} \mathfrak{z}_t = 1$ for any $t \in [0, T]$, it suffices to show the family $\{\mathfrak{z}_T^n\}_{n \rightarrow \infty}$ is uniformly integrable. With chosen σ_n , we have $\mathbb{E} \mathfrak{z}_T^n = 1$. Let us introduce a probability measure $\tilde{\mathbb{P}}_T^n \ll \mathbb{P}$ with $d\tilde{\mathbb{P}}_T^n = \mathfrak{z}_T^n d\mathbb{P}$ and denote by $\tilde{\mathbb{E}}_T^n$ the expectation symbol of $\tilde{\mathbb{P}}_T^n$.

Following Hitsuda [4], the uniform integrability of $\{\mathfrak{z}_T^n\}_{n \rightarrow \infty}$ is verified with a convex function

$$\psi(x) = x \log(x) + 1 - x, \quad x \geq 0 \quad (1.4)$$

due to the Vallée-Poussin's criteria since $\lim_{x \rightarrow \infty} \frac{\psi(x)}{x} = \infty$. Namely, we have to show that $\sup_n \mathbb{E} \psi(\mathfrak{z}_T^n) < \infty$. A verification of this condition is inconvenient. However if $\mathbb{E} \psi(\mathfrak{z}_T^n) < \infty$ a direct computation shows $\mathbb{E} \psi(\mathfrak{z}_T^n) = \tilde{\mathbb{E}}_T^n \log(\mathfrak{z}_T^n)$.

(3) the Girsanov theorem, a random process $(\tilde{B}_t^n)_{t \in [0, T]}$ with

$$\tilde{B}_t^n = B_t^n - \int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) ds \quad (1.5)$$

is $\tilde{\mathbb{P}}_T^n$ -martingale with the predictable quadratic variation $\langle \tilde{B}^n \rangle_t = \langle B^n \rangle_t \equiv t \wedge \sigma_n$. Therefore, by using (1.5), we obtain

$$\log(\mathfrak{z}_t^n) \stackrel{\tilde{\mathbb{P}}_T^n}{=} \int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) d\tilde{B}_s^n + \frac{1}{2} \int_0^t I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds.$$

Since $I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s)$ is bounded, we have $\tilde{\mathbb{E}}_T^n \int_0^T I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds \leq c_n$ with a constant c_n depending on n . Hence, $\tilde{\mathbb{E}}_T^n \int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) d\tilde{B}_s^n = 0$.

Therefore and in view of $I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds \leq \text{const.} [1 + \sup_{s' \in [0, s \wedge \sigma_n]} |B_{s'}^n|^2]$,

$$\tilde{\mathbb{E}}_T^n \log(\mathfrak{z}_T^n) = \tilde{\mathbb{E}}_T^n \int_0^T I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds \leq \text{const.} \left[1 + \tilde{\mathbb{E}}_T^n \sup_{s' \in [0, T \wedge \sigma_n]} |B_{s'}^n|^2 \right].$$

(4) Now, the proof is reduced to $\sup_n \tilde{\mathbb{E}}_T^n \sup_{s' \in [0, T \wedge \sigma_n]} |B_{s'}^n|^2 < \infty$.

Denote $V_t^n := \tilde{\mathbb{E}}_T^n \sup_{s' \in [0, t \wedge \sigma_n]} |B_{s'}^n|^2$. In view of (1.5),

$$V_t^n \leq 2\tilde{\mathbb{E}}_T^n \left(\int_0^t I_{\{\sigma_n \geq s\}} |\alpha(\omega, s)| ds \right)^2 + 2\tilde{\mathbb{E}}_T^n \left(\sup_{s' \in [0, t \wedge \sigma_n]} |\tilde{B}_{s'}^n| \right)^2.$$

By the Doob maximal inequality, $\tilde{\mathbb{E}}_T^n \left(\sup_{s' \in [0, t \wedge \sigma_n]} |\tilde{B}_{s'}^n| \right)^2 \leq 4\tilde{\mathbb{E}}_T^n |\tilde{B}_{t \wedge \sigma_n}^n|^2 = 4\tilde{\mathbb{E}}_T^n |(t \wedge \sigma_n)| \leq 4T$, while by the Cauchy-Schwarz inequality

$$\tilde{\mathbb{E}}_T^n \left(\int_0^t I_{\{\sigma_n \geq s\}} |\alpha(\omega, s)| ds \right)^2 \leq \tilde{\mathbb{E}}_T^n \int_0^t I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds \leq \mathbf{r} \left[1 + \int_0^t V_s^n ds \right].$$

Finally, combining these estimates, we obtain an integral inequality (with the constant $\mathbf{r}$ independent of n):

$$V_t^n \leq \mathbf{r} \left[1 + \int_0^t V_s^n ds \right].$$

Thus, $V_T^n \leq \mathbf{r} e^{T\mathbf{r}}$.

1.3. Formulation of main result. Let us explain how Brownian B_t might be replaced by a purely discontinuous martingale M_t . A simplest way is to replace B_t by M_t in the Doléans-Dade equation (1.3), that is,

$$\mathfrak{z}_t = 1 + \int_0^t \mathfrak{z}_{s-} \alpha(\omega, s) dM_s,$$

where $\mathfrak{z}_{s-} = \lim_{s' \uparrow s} \mathfrak{z}_{s'}$, and adapted process $\alpha(\omega, t)$ is replaced by its predictable version. The square integrable martingale M_t has paths the Skorokhod space $\mathbb{D}_{[0, \infty)}$. Denote $\langle M \rangle_t$ the predictable quadratic variation of M_t and $M_{t-} = \lim_{t' \uparrow t} M_{t'}$.

We assume $\int_0^t \alpha^2(\omega, s) d\langle M \rangle_s < \infty$. A positiveness of $\mathfrak{z}_t$ is warranted by assumptions $\alpha(\omega, t) \geq 0$ and $(M_t - M_{t-}) I_{\{M_t \neq M_{t-}\}} > 0$.

We choose M_t in a form of Itô's integral

$$M_t = \int_0^t \int_{\mathbb{R}_+} z [\mu(ds, dz) - \nu(ds, dz)]$$

relative to, so called, martingale difference “ $\mu - \nu$ ”, where $\mu(dt, dz)$ is the integer-valued measure $\mu = \mu(dt, dz)$ associated with a jump process $\triangle M_t = M_t - M_{t-}$ of M_t and $\nu(dt, dz)$ is a compensator of $\mu(dt, dz)$. In order to have the above-mentioned properties of M_t , we choose a deterministic compensator

$$\nu(dt, dz) = K(dz)dt$$

with a measure $K(dz)$ supported on $\mathbb{R}_+$ and $\int_{\mathbb{R}_+} z^2 K(dz) < \infty$. In particular, then,

$$\langle M \rangle_t \equiv \mathbb{E} M_t^2 \equiv \int_0^t z^2 K(dz) ds.$$

Our main result is formulated in

Theorem 1.1. Assume $\int_{\mathbb{R}_+} z^3 K(dz) < \infty$. Then for any $T > 0$ (comp. (1.2))

$$|\alpha(\omega, t)|^2 \leq \text{const.} \left[1 + \sup_{s \in [0, t]} M_{s-}^2 \right]_{t \in [0, T]} \Rightarrow \mathbf{E} \mathfrak{z}_T = 1,$$

The method of proof is similar to one given in Section (1.2)

2. The proof of Theorem 1.1

2.1. Preliminaries. We begin with recalling necessary notions (for more details, see e.g. [8] or [3]). Along this paper a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, \infty)}, \mathbf{P})$ with “general conditions” is fixed and all random objects are defined on it. $\mathcal{P}$ denotes predictable σ -algebra relative to $(\mathcal{F}_t)_{t \in [0, \infty)}$ and $\mathcal{B}_+$ is the Borel σ -algebra on $\mathbb{R}_+$.

Henceforth, $\mathbf{r}$ denotes a generic constant taking different values at different appearances and is independent of a number n involved in the text.

We begin with know implication:

$$|\alpha(\omega, t)| \leq \mathbf{r} \Rightarrow \mathbf{E} \mathfrak{z}_t \equiv 1. \quad (2.1)$$

Set $\tau_n = \inf\{t : \mathfrak{z}_t \geq n\}$ and notice $\mathfrak{z}_{(t \wedge \tau_n)-} \leq n$. Then $\mathfrak{z}_t^n := \mathfrak{z}_{t \wedge \tau_n}$ solves the Doleans-Dade equation

$$\mathfrak{z}_t^n = 1 + \int_0^t \mathfrak{z}_{s-}^n I_{\{\tau_n \geq s\}} \alpha(\omega, s) dM_s. \quad (2.2)$$

A boundedness of $I_{\{\tau_n \geq s\}} \mathfrak{z}_{s-}^n \alpha(\omega, s)$ guarantees the process $\mathfrak{z}_t^n$ is the square integrable martingale and $\mathbf{E}(\mathfrak{z}_t^n)^2 = 1 + \mathbf{E} \int_0^t (I_{\{\tau_n \geq s\}} \mathfrak{z}_{s-}^n \alpha(\omega, s))^2 d\langle M \rangle_s ds \leq 1 + \mathbf{r} \int_0^t \mathbf{E}(\mathfrak{z}_s^n)^2 ds$. Then, a function $V_t^n = \mathbf{E}(\mathfrak{z}_t^n)^2$ solves the integral inequality: $V_t^n \leq 1 + \mathbf{r} \int_0^t V_s^n ds$. So, by the Bellman-Gronwall inequality, $V_t^n \leq e^{\mathbf{r}t}$, that is, $\sup_n \mathbf{E}(\mathfrak{z}_t^n)^2 \leq e^{\mathbf{r}t}$. Therefore, by the Vallée-Poussin's criteria, the family $\{\mathfrak{z}_t^n\}_{n \rightarrow \infty}$ is uniformly integrable. So, not only $\lim_{n \rightarrow \infty} \mathfrak{z}_t^n = \mathfrak{z}_t$ but also $\mathbf{E} \mathfrak{z}_t = \lim_{n \rightarrow \infty} \mathbf{E} \mathfrak{z}_t^n \equiv 1$.

2.2. $\mathfrak{z}_t^n$ approximation of $\mathfrak{z}_t$. Change of probability measure.

Lemma 2.1. Let $\sigma_n = \inf\{t : [1 + \sup_{s \in [0, t]} M_{s-}^2] \geq n\}$ and $\mathfrak{z}_t^n = \mathfrak{z}_{t \wedge \sigma_n}$. Then $\mathbf{E} \mathfrak{z}_t^n \equiv 1$.

Proof. We use (2.2) with τ_n replaced by σ_n :

$$\mathfrak{z}_t^n = 1 + \int_0^t I_{\{\sigma_n \geq s\}} \mathfrak{z}_{s-}^n \alpha(\omega, s) dM_s. \quad (2.3)$$

Since $[1 + \sup_{s \in [0, t \wedge \sigma_n]} M_{s-}^2] \leq n$ and, then, $I_{\{\sigma_n \geq t\}} |\alpha(\omega, t)|^2 \leq \mathbf{r} I_{\{\sigma_n \geq t\}} [1 + \sup_{s \in [0, t]} M_{s-}^2] \leq \mathbf{r} n$, it remains to apply (2.1). $\square$

Let $T > 0$ be fixed. By Lemma 2.1 $\mathbf{E} \mathfrak{z}_T^n = 1$. As in Section 1.2, we have to show the family $\{\mathfrak{z}_T^n\}_{n \rightarrow \infty}$ is uniformly integrable. So, we intend to verify $\sup_n \mathbf{E} \psi(\mathfrak{z}_T^n) < \infty$ with the function $\psi(x)$ defined in (1.4).

Repeating arguments from Section 1.2, we introduce a probability measure $\tilde{\mathbf{P}}_T^n \ll \mathbf{P}$ with $d\tilde{\mathbf{P}}_T^n = \mathfrak{z}_T^n d\mathbf{P}$ ($\tilde{\mathbf{E}}_T^n$ denotes the expectation symbol of $\tilde{\mathbf{P}}_T^n$). Set $M_t^n = M_{t \wedge \sigma_n}$ and rewrite (2.3) in equivalent form

$$\mathfrak{z}_t^n = 1 + \int_0^t I_{\{\sigma_n \geq s\}} \mathfrak{z}_{s-}^n \alpha(\omega, s) dM_s^n. \quad (2.4)$$

The random process $(M_t^n)_{t \in [0, T]}$ is $\mathbf{P}$ - square integrable martingale. Since $\tilde{\mathbf{P}}_T^n \ll \mathbf{P}$, the process $(M_t^n)_{t \in [0, T]}$ is $\tilde{\mathbf{P}}_T^n$ -semimartingale obeying the unique decomposition $M_t^n = A_t^n + \tilde{M}_t^n$ with predictable drift A_t^n and local martingale $\tilde{M}_t^n$ (see, e.g. [8], Ch.4, §5, Theorem 2). Denote $\tilde{\nu}_T^n(dt, dz)$ a compensator of $\mu(dt, dz)$ relative to $\tilde{\mathbf{P}}_T^n$.

Lemma 2.2.

1. $\tilde{\nu}_T^n(ds, dz) = I_{\{\sigma_n \geq s\}}(1 + \alpha(\omega, s)z)K(dz)ds$
2. $\tilde{M}_t^n = M_t^n - A_t^n$ is square integrable martingale with

$$\langle \tilde{M}^n \rangle_t = \int_0^t \int_{\mathbb{R}_+} z^2 I_{\{\sigma_n \geq s\}}(1 + \alpha(\omega, s)z)K(dz)ds \quad (2.5)$$

3. $A_t^n = \int_0^t \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} \alpha(\omega, s) z^2 K(dz)ds$

Proof. 1. Below, we will use a formula

$$\frac{\mathfrak{z}_s^n}{\mathfrak{z}_{s-}^n} I_{\{\sigma_n \geq s\}} = I_{\{\sigma_n \geq s\}}(1 + \alpha(\omega, s) \triangle M_s^n)$$

readily derived from (2.4). Let $u(\omega, t, z)$ be bounded and $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}_+)$ -measurable function vanishing in a vicinity of $\{0\}$. Write

$$\begin{aligned} & \tilde{\mathbf{E}}_T^n \int_0^T \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} u(\omega, s, z) \mu(dt, dz) \\ &= \mathbf{E} \mathfrak{z}_T^n \int_0^T \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} u(\omega, s, z) \mu(dt, dz) \\ &= \mathbf{E} \int_0^T \int_{\mathbb{R}_+} \mathfrak{z}_s^n I_{\{\sigma_n \geq s\}} u(\omega, s, z) \mu(dt, dz) \\ &= \mathbf{E} \int_0^T \int_{\mathbb{R}_+} \mathfrak{z}_{s-}^n \frac{\mathfrak{z}_s^n}{\mathfrak{z}_{s-}^n} I_{\{\sigma_n \geq s\}} u(\omega, s, z) \mu(dt, dz) \\ &= \mathbf{E} \int_0^T \int_{\mathbb{R}_+} \mathfrak{z}_{s-}^n [1 + \alpha(\omega, s) \triangle M_s^n] I_{\{\sigma_n \geq s\}} u(\omega, s, z) \mu(dt, dz) \\ &= \mathbf{E} \int_0^T \int_{\mathbb{R}_+} \mathfrak{z}_{s-}^n I_{\{\sigma_n \geq s\}} [1 + \alpha(\omega, s)z] u(\omega, s, z) \mu(ds, dz) \\ &= \mathbf{E} \int_0^T \int_{\mathbb{R}_+} \mathfrak{z}_{s-}^n I_{\{\sigma_n \geq s\}} [1 + \alpha(\omega, s)z] u(\omega, s, z) K(dz)ds \\ &= \mathbf{E} \mathfrak{z}_T^n \int_0^T \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} [1 + \alpha(\omega, s)z] u(\omega, s, z) K(dz)ds \\ &= \tilde{\mathbf{E}}_T^n \int_0^T \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} [1 + \alpha(\omega, s)z] u(\omega, s, z) K(dz)ds. \end{aligned}$$

The desired result follows by arbitrariness of $u(\omega, t, z)$.

2.+3. Recall $\int_{\mathbb{R}_+} (z^2 \vee z^3) K(dz) < \infty$. Hence $\tilde{\mathbf{E}}_T^n \int_0^T \int_{\mathbb{R}_+} z^2 \tilde{\nu}^n(ds, dz) < \infty$. Therefore

$$\widehat{M}_t^n = \int_0^t \int_{\mathbb{R}_+} z I_{\{\sigma_n \geq s\}} [\mu(ds, dz) - \tilde{\nu}_T^n(ds, dz)]$$

is $\tilde{\mathbf{P}}_T^n$ -square integrable martingale with predictable quadratic variation process

$$\langle \widehat{M}^n \rangle_t = \int_0^t \int_{\mathbb{R}_+} z^2 I_{\{\sigma_n \geq s\}} (1 + \alpha(\omega, s)z) K(dz) ds. \quad (2.6)$$

On the other hand,

$$\begin{aligned} M_t^n - \widehat{M}_t^n &= \int_0^t \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} z [\tilde{\nu}_T^n(ds, dz) - K(dz)] ds \\ &= \int_0^t \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} z^2 \alpha(\omega, s) K(dz) ds =: A_t^n, \end{aligned}$$

where A_t^n is well defined predictable process.

Consequently, $\widehat{M}_t^n \equiv \widetilde{M}_t^n$ in view of the unique semimartingale decomposition with the predictable drift. $\square$

2.3. Upper bound of $\mathbf{E}\psi(\mathfrak{z}_T^n)$. Following the main idea of Section 1.2, we have to show the family $\{\mathfrak{z}_T^n\}_{n \rightarrow \infty}$ is uniformly integrable. To this end, we have to prove $\sup_n \mathbf{E}\psi(\mathfrak{z}_T^n) < \infty$ for the function $\psi(x) = x \log(x) + 1 - x$, $x \geq 0$. We show first

$$\tilde{\mathbf{E}}_T^n \log(\mathfrak{z}_T^n) < \infty$$

and, then, use an obvious identity $\mathbf{E}\psi(\mathfrak{z}_T^n) = \tilde{\mathbf{E}}_T^n \log(\mathfrak{z}_T^n)$.

Lemma 2.3.

$$\tilde{\mathbf{E}}_T^n \log(\mathfrak{z}_T^n) \leq \tilde{\mathbf{E}}_T^n \int_0^T \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} z^2 \alpha^2(\omega, s) K(dz) ds \leq \mathbf{r} \left[1 + \tilde{\mathbf{E}}_T^n \sup_{s \in [0, T \wedge \sigma_n]} (M_{s-}^n)^2 \right].$$

Proof. It is well known (see e.g. [8], Ch. 2, §4), the Doleans-Dade equation (2.4) obeys the unique solution

$$\begin{aligned} \mathfrak{z}_t^n &= \exp \left(\int_0^t \int_{\mathbb{R}_+} I_{\{\sigma_n \geq s\}} \alpha(\omega, s) dM_s^n \right. \\ &\quad \left. + \sum_{s \in [0, t \wedge \sigma_n]} \log \left\{ [1 + \alpha(\omega, s) \Delta M_s^n] - \alpha(\omega, s) \Delta M_s^n \right\} \right). \end{aligned}$$

Recall $\alpha(\omega, s) \Delta M_s^n \geq 0$. Then $\log[1 + \alpha(\omega, s) \Delta M_s^n] - \alpha(\omega, s) \Delta M_s^n \leq 0$ and, therefore, $\log(\mathfrak{z}_t^n) \leq \int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) dM_s^n$. So, by Lemma 2.2,

$$\log(\mathfrak{z}_T^n) \leq \int_0^T I_{\{\sigma_n \geq s\}} \alpha(\omega, s) dA_s^n + \int_0^T I_{\{\sigma_n \geq s\}} \alpha(\omega, s) d\widetilde{M}_s^n.$$

The process $\widetilde{M}_t^n$ is square integrable martingale with $\langle \widetilde{M}^n \rangle_T$ defined in (2.5). Therefore, the Itô integral $\int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) d\widetilde{M}_s^n$ is also the square integrable martingale with the quadratic variation

$$QV := \int_0^T \int_{\mathbb{R}_+} z^2 I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) [1 + \alpha(\omega, s)z] K(dz) ds.$$

Let us show that QV is bounded by a constant depending on n . Since $\int_{\mathbb{R}_+} (z^2 \vee z^3) K(dz) \leq \mathbf{r}$,

$$QV \leq \mathbf{r} \int_0^T I_{\{\sigma_n \geq s\}} [1 + \alpha^3(\omega, s)] ds \leq \mathbf{r}T + \mathbf{r} \int_0^T I_{\{\sigma_n \geq s\}} \alpha^3(\omega, s) ds.$$

Further,

$$I_{\{\sigma_n \geq s\}} \alpha^3(\omega, s) = [I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s)]^{3/2} \leq \mathbf{r} \left[1 + \sup_{s' \in [0, s \wedge \sigma_n]} M_{s'-}^2 \right]^{3/2} \leq \mathbf{r} n^{3/2},$$

that is, $QV \leq \mathbf{r}T[1 + n^{3/2}]$.

Hence, $\widetilde{E}_T^n \int_0^T I_{\{\sigma_n \geq s\}} \alpha(\omega, s) d\widetilde{M}_s^n = 0$ and $\widetilde{E}_T^n \log(\mathfrak{z}_T^n) \leq \widetilde{E}_T^n \int_0^T I_{\{\sigma_n \geq s\}} \alpha(\omega, s) dA_s^n$. So, it remains to recall the formula of A_t^n (see Lemma 2.2), and $\int_{\mathbb{R}_+} z^2 K(dz)$, and

$$I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) = I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) \leq \mathbf{r} \left[1 + \sup_{s' \in [0, s \wedge \sigma_n]} M_{s'-}^2 \right].$$

□

2.4. Final step of the proof. Now, we are in the position to compute $\widetilde{E}_T^n \sup_{s \in [0, T]} |M_s^n|^2$.

The use of $M_t^n = A_t^n + \widetilde{M}_t^n$ implies $\widetilde{E}_T^n \sup_{t' \in [0, t]} |M_{t'}^n|^2 \leq 2\widetilde{E}_T^n \sup_{t' \in [0, t]} |A_{t'}^n|^2 + 2\widetilde{E}_T^n \sup_{t' \in [0, t]} |\widetilde{M}_{t'}^n|^2$.

In view of statement **3.** of Lemma 2.2 $A_t^n = \int_{\mathbb{R}_+} z^2 K(dz) \int_0^t I_{\{\sigma_n \geq s\}} \alpha(\omega, s) ds$. So, by applying the Cauchy-Schwarz inequality we obtain

$$\widetilde{E}_T^n \sup_{t' \in [0, t]} |A_{t'}^n|^2 \leq \mathbf{r} \widetilde{E}_T^n \int_0^t I_{\{\sigma_n \geq s\}} \alpha^2(\omega, s) ds. \quad (2.7)$$

Further, by the Doob maximal inequality and (2.6) we find that

$$\begin{aligned} \widetilde{E}_T^n \sup_{t' \in [0, t]} |\widetilde{M}_{t'}^n|^2 &\leq 4\widetilde{E}_T^n \langle \widetilde{M}^n \rangle_t = 4\widetilde{E}_T^n \int_0^t \int_{\mathbb{R}_+} z^2 I_{\{\sigma_n \geq s\}} (1 + \alpha(\omega, s)z) K(dz) ds \\ &\leq \mathbf{r} \widetilde{E}_T^n \int_0^t I_{\{\sigma_n \geq s\}} [1 + \alpha(\omega, s)] ds \leq \mathbf{r} \widetilde{E}_T^n \int_0^t I_{\{\sigma_n \geq s\}} [1 + \alpha^2(\omega, s)] ds \end{aligned} \quad (2.8)$$

Now, a combination of (2.7), and (2.8) provides: for any $t \leq T$,

$$\widetilde{E}_T^n \sup_{t' \in [0, t]} |M_{t'}^n|^2 \leq \mathbf{r} \widetilde{E}_T^n \int_0^t [1 + \alpha^2(\omega, s)] ds \leq \mathbf{r} \widetilde{E}_T^n \int_0^t \left[1 + \sup_{s' \in [0, s]} |M_{s'}^n|^2 \right] ds.$$

Hence, the function $V_t^n := \sup_n \widetilde{E}_T^n \sup_{s \in [0, t]} |M_s^n|^2$ solves an integral inequality:

$$“V_t^n \leq \mathbf{r} \left(1 + \int_0^t V_s^n ds \right)” \Rightarrow “V_T^n \leq \mathbf{r} e^{\mathbf{r}T}”.$$



REFERENCES

- [1] Benes, V.E. (1971) Existence of optimal stochastic control laws *SIAM J. of Control*, 9 , 446-475
- [2] Girsanov, I.V. (1960) On transforming a certain class of stochastic processes by absolutely continuous substitution of measures. *Theory Probab. Appl.* **5**, 285-301.
- [3] Jacod J., Shiryaev A.N.: Limit theorems for stochastic processes. 2nd ed. Springer-Verlag, Berlin (2003)
- [4] Hitsuda Masuyuki. (1968) Representation of Gaussian processes equivalent to Wiener process. *Osaka J. Math.* **5**, 299-312.
- [5] Karatzas, I. and Shreve, S.E. (1991): Brownian Motion and Stochastic Calculus. Springer-Verlag, New York Berlin Heidelberg.
- [6] Kazamaki, N. (1977) On a problem of Girsanov.// Tôhoku Math. J., 29 , p. 597- 600.
- [7] Krylov, N.V. (8 May 2009) A simple proof of a result of A. Novikov. arXiv:math/020713v2 [math.PR]
- [8] Liptser, R.Sh., Shiryaev, A.N. (1989) *Theory of Martingales*. Kluwer Acad. Publ.
- [9] Liptser, R. Sh. and Shiryaev, A. N. (2000). *Statistics of Random Processes I*, 2nd ed., Springer, Berlin - New York.
- [10] Novikov, A.A. (1979) On the conditions of the uniform integrability of the continuous nonnegative martingales.// Theory of Probability and its Applications, 24, No. 4, p. 821-825.
- [11] "Ustünel. A and Zakai, M. (2000) *Transformation of measure on Wiener space*. Springer, Berlin - New York.

DEPARTMENT OF ELECTRICAL ENGINEERING SYSTEMS, TEL AVIV UNIVERSITY, 69978 TEL AVIV, ISRAEL
E-mail address: `liptser@eng.tau.ac.il`; `rliptser@gmail.com`